



## Research report

# Behavioral deficits induced by third-trimester equivalent alcohol exposure in male C57BL/6J mice are not associated with reduced adult hippocampal neurogenesis but are still rescued with voluntary exercise



G.F. Hamilton (PhD)\*, P.J. Bucko (BS), D.S. Miller (BS), R.S. DeAngelis (BA), C.P. Krebs (BS), J.S. Rhodes (PhD)

*Department of Psychology, The Beckman Institute, 405 N Mathews Ave, University of Illinois at Urbana-Champaign, Urbana, IL, 61801, USA*

## HIGHLIGHTS

- PD5, 7 and 9 but not PD5 alcohol exposure impairs passive avoidance acquisition.
- PD5, 7 and 9 but not PD5 alcohol exposure impairs rotarod performance.
- Neonatal alcohol exposure does not impact adult neurogenesis levels.

## ARTICLE INFO

*Article history:*

Received 6 May 2016

Received in revised form 21 July 2016

Accepted 30 July 2016

Available online 1 August 2016

*Keywords:*

Bromodeoxyuridine

Passive avoidance

Running

Hippocampus

Fetal alcohol spectrum disorders

## ABSTRACT

Prenatal alcohol exposure can produce permanent alterations in brain structure and profound behavioral deficits. Mouse models can help discover mechanisms and identify potentially useful interventions. This study examined long-term influences of either a single or repeated alcohol exposure during the third-trimester equivalent on survival of new neurons in the hippocampus, behavioral performance on the Passive avoidance and Rotarod tasks, and the potential role of exercise as a therapeutic intervention. C57BL/6J male mice received either saline or 5 g/kg ethanol split into two s.c. injections, two hours apart, on postnatal day (PD)7 (Experiment 1) or on PD5, 7 and 9 (Experiment 2). All mice were weaned on PD21 and received either a running wheel or remained sedentary from PD35–PD80/81. From PD36–45, mice received i.p. injections of 50 mg/kg bromodeoxyuridine (BrdU) to label dividing cells. Behavioral testing occurred between PD72–79. Number of surviving BrdU+ cells and immature neurons (doublecortin; DCX+) was measured at PD80–81. Alcohol did not affect number of BrdU+ or DCX+ cells in either experiment. Running significantly increased number of BrdU+ and DCX+ cells in both treatment groups. Alcohol-induced deficits on Rotarod performance and acquisition of the Passive avoidance task (Day 1) were evident only in Experiment 2 and running rescued these deficits. These data suggest neonatal alcohol exposure does not result in long-term impairments in adult hippocampal neurogenesis in the mouse model. Three doses of ethanol were necessary to induce behavioral deficits. Finally, the mechanisms by which exercise ameliorated the neonatal alcohol induced behavioral deficits remain unknown.

© 2016 Elsevier B.V. All rights reserved.

## 1. Introduction

Approximately 2–5% of all live births in the US are diagnosed as cases of Fetal Alcohol Spectrum Disorders (FASD), an unchanging number despite clear knowledge and the preventable nature of

FASD [48]. In humans, prenatal alcohol exposure produces a wide range of long-lasting neurobehavioral deficits, including physical, cognitive, learning and behavioral disabilities [7]. For example, FASD patients exhibit long-lasting deficits in working memory and behavioral flexibility [14,58], behaviors that require the proper functioning of multiple brain regions at once, many of which are still developing during and appear particularly sensitive to alcohol exposure during the third-trimester.

The third trimester is comparable to the first two weeks of postnatal life in mice and rats in terms of brain development [17]. A

\* Corresponding author at: 3315 Beckman Institute, 405 North Mathews Avenue, Urbana, Illinois 61801, USA.

E-mail address: [gillianh@illinois.edu](mailto:gillianh@illinois.edu) (G.F. Hamilton).


large literature has established that rodents neonatally exposed to alcohol exhibit impaired performance on a variety of behavioral tasks as adults (for review see [54]. Of note, only a few studies have demonstrated a significant influence of neonatal alcohol exposure on Passive avoidance performance [1,3,4,61], a task in which mice must learn and remember to inhibit their natural response to move to the dark side of the chamber and instead stay in the lit side of the chamber. This task involves multiple brain regions [38,69], including, but not limited to, the prefrontal cortex and the hippocampus, two brain regions well-known for being particularly sensitive to third-trimester equivalent alcohol exposure. In fact, Barron and colleagues [3], using an artificial rearing model, demonstrated that third-trimester equivalent alcohol exposure impaired both acquisition and retention of this task in PD23 rats. Whether a similar result is evident in mice and whether this effect persists into adulthood remains unknown. However, given the large number of adult FASD patients suffering from working memory and behavioral flexibility deficits, this is a well-suited behavioral task to use when developing interventions.

Neonatal alcohol exposure produces long-lasting neuroanatomical deficits in the hippocampus of rodents. This is particularly apparent following a third-trimester equivalent exposure [20,46]. Of all the brain regions impacted from alcohol, the hippocampus is arguably the most capable of extensive regeneration and repair [6,47,53]. Still, it is possible that neonatal exposure can produce a persistent reduction in hippocampal plasticity. Some studies show reductions in adult hippocampal neurogenesis and suggest the loss contributes to behavioral deficits [22,27,34,42], while, other studies fail to find an effect of neonatal ethanol exposure on adult neurogenesis [9,22,30,71]. These studies used both rats and mice of different strains. The lack of consensus warrants additional research specifically examining the impact of different timing and duration of alcohol exposures on adult hippocampal neurogenesis in a single mouse strain.

Regardless of whether or not neonatal alcohol exposure results in long-lasting deficits in adult neurogenesis in rodents, exercise is known to increase levels of adult neurogenesis and improve behavioral performance across a broad range of tasks in alcohol-exposed rats [10,21,24,26,60]. Therefore, increasing levels of adult neurogenesis in alcohol-exposed mice could rescue hippocampal function either through compensation or repair. The goal of this study was to determine the extent to which a single (PD7) or repeated (PD5, 7 and 9) binge ethanol exposure reduces adult hippocampal neurogenesis, and impairs behavioral performance on the Rotarod and Passive avoidance tasks, and whether exercise can mitigate any deficits.

## 2. Materials and methods

Seventy-one (Experiment 1:36; Experiment 2:35) male C57BL/6J mice (Jackson Laboratory, Bar Harbor, ME) as adults were used plus an additional 37 pups euthanized at PD7 for blood-ethanol concentration measurements. The 71 mice were generated from 20 pairs (10 pairs per Experiment), and the 37 pups from 7 pairs. The 71 mice were randomized into four groups: saline-sedentary, saline-runner, alcohol-sedentary, alcohol-runner (Experiment 1: n = 9 per group; Experiment 2, n = 7, 8, 10, and 10, respectively). No more than two animals per litter were assigned to a given treatment condition. Mice were housed in standard polycarbonate shoebox cages (29 cm x 19 cm x 13 cm; L x W x H) with corncob bedding (Teklad 7012; Harlan Teklad, Madison, WI, USA). Rooms were controlled for temperature (21 °C ± 1 °C) and photo-period (12:12 D:L; lights off at 10:00am and on at 10:00pm). Food and water was provided *ad libitum*. The Beckman Institute Animal Facility is AAALAC approved. All procedures were



**Fig. 1.** Experimental Timeline. For Experiment 1, mice were exposed to a single day of treatment, of either sterile saline or 5 g/kg ethanol (split into two doses, two hours apart), on PD 7. For Experiment 2, mice received three days of treatment (sterile saline or 5 g/kg ethanol) on PD 5, 7, and 9. All mice were weaned on PD 21 and singly housed on PD 35, the onset of intervention, which consisted of either voluntary access to a running wheel or sedentary condition. Newly dividing cells were labeled with bromodeoxyuridine (BrdU) for the first ten days of intervention. Mice underwent Passive avoidance and Rotarod testing during adulthood, after being exposed to the assigned intervention for at least thirty days. Differences in the order of the testing procedures were due to the availability of testing equipment.

approved by the University of Illinois Institutional Animal Care and Use Committee and adhered to NIH guidelines. All efforts were made to minimize the number of mice used and their suffering.

### 2.1. Neonatal treatment

**Experiment 1:** We followed established methods to deliver ethanol to neonatal mice [34,43,71]. Pups from ten litters were injected subcutaneously (s.c.) over the shoulders, into the loose skin over the neck, with either a sterile saline solution or a dose of 5 g/kg of ethanol (between 0.03 – 0.08 mL of a 20% ethanol solution per injection; Decon Laboratories, Inc., King of Prussia, PA, USA; 20% ethanol in sterile saline solution; 15.8 mL/kg), which was divided into two injections, two hours apart. To avoid leakage of the solution, the liquid was released slowly into the animal and the needle remained in the animal for 3–5 s following before slowly being removed in order to allow the liquid time to be absorbed. Whole litters were injected with either saline or ethanol. The males from each litter were then assigned to the 2 treatments (runner or sedentary). Litters had 2–4 males, hence there were 1 or 2 males from the same litter per treatment combination. Dose was chosen based on the literature [34,35,43,68,71]. On PD21, all mice were weaned into groups of four by sex (Fig. 1).

**Experiment 2:** The same procedure as described above was used with the exception that mice received treatments on PD5, 7, and 9.

### 2.2. Blood ethanol concentration analysis

Trunk blood samples were collected from an additional cohort of 37 mice in order to verify that alcohol concentrations in the blood (BECs) reached a toxic level, above 200 mg/dl for four consecutive hours or more, as BECs at this level have been shown to produce widespread apoptotic neurodegeneration [35]. Mice were assigned to one of six groups at PD7: control (no alcohol administered; n = 3) or alcohol administered then trunk blood collected at the following time points after the first injection of alcohol: 30 min (n = 8), 1 h (n = 8), 3 h (n = 7), 5 h (n = 5), and 9 h (n = 6). Note that for all time points greater than or equal to 3 h, the mice would have received a second injection at 2 h for a total of 5 g/kg. Immediately following collection in heparinized tubes, trunk blood was centrifuged at

1000 × g at 4 °C for 10 min after which plasma was collected and immediately stored at –80 °C.

Plasma samples were analyzed using the Colorimetric Alcohol Assay Kit (STA-620; Cell Biolabs, Inc, San Diego, CA, USA). Briefly, the stock Ethanol Standard was diluted using deionized water in order to prepare an ethanol standard (200 mM). Said ethanol standard was then further diluted in a 1X Assay Buffer to produce standards in the concentration range of 0–200 µm. Next, all samples were diluted in 1X Assay Buffer (1:80). Following this, 10 µL of the diluted ethanol standards or samples were added to a 96-well microtiter plate (all standards and samples were run in duplicate). The microtiter plate was then covered and placed to the side while the Reaction Mixture was prepared. The Reaction Mixture consisted of 8.7 mL of deionized water, 1 mL of 10X Assay Buffer, 100 µL of 100X Enzyme Mixture and 200 µL of the 50X Colorimetric Probe. The Reaction Mixture was mixed and then 90 µL was added to each well in the microtiter plate. The plate was covered and subsequently incubated at 37 °C for 30 min. Finally, absorbances were read at 570 nm on a microplate reader (EPOCH; Bitek Instruments Inc, Winooski, VT, USA) and the ethanol concentration of samples was calculated.

### 2.3. Exercise or sedentary intervention

On PD35, all mice (Experiment 1 and Experiment 2) were singly housed with or without running wheels in order to obtain individual running levels. The intervention was counterbalanced within the treatments. Sedentary mice were housed in standard polycarbonate shoebox cages, as previously described. Running mice were housed in cages (36 cm × 20 cm × 14 cm; L × W × H) with a 23 cm diameter wheel mounted to the cage top (Respironics, Bend, OR, USA). Wheel rotations were recorded continuously in one minute increments via magnetic switches interfaced to a computer throughout the experiment. Mice assigned to the sedentary condition were not housed in cages with locked wheels because mice climb in locked wheels and we intended to keep physical activity to a minimum in the sedentary group. Similar to previous studies from our lab, mice were given at least 30 days access to the exercise intervention prior to any behavioral intervention.

### 2.4. 5'bromo-5'-deoxyuridine (BrdU) Injections

Following established procedures, from PD36–45, all mice received a daily intraperitoneal (*i.p.*) injection of BrdU (50 mg/kg) to label newly dividing cells in the dentate gyrus of the hippocampus [9,30,34,42].

### 2.5. Rotarod

Given that children with FASD typically have poor motor coordination, we wanted to determine first whether a PD7 or a PD5,7 and 9 alcohol exposure could impair motor coordination, and second, whether exercise could ameliorate any alcohol-induced deficit. Mice underwent Rotarod testing beginning one hour after lights off either on PD76–77 (Experiment 1) or PD75–76 (Experiment 2). Mice were placed on a stationary Rotarod (AccuRotor Rota Rod Tall Unit, 63 cm fall height, 30 mm diameter rotating dowel; Accuscan, Columbus, OH, USA), the dowel was accelerated at 60 rpm/min, and latency to fall recorded. Mice were tested for two consecutive days with four consecutive trials per day.

### 2.6. Passive avoidance

Experiment 1: Mice underwent Passive avoidance testing on PD78–79 beginning one hour after lights off. On PD78, mice were trained in the Passive avoidance chamber (San Diego Instruments,

San Diego, CA, USA). The chamber consisted of two compartments, one lit and one unlit, separated by a guillotine door. The bottom of the chamber contained a metal grid through which a mild foot shock could be delivered. Each mouse was placed in the lit side of a chamber and the guillotine door was raised. Latency to cross over into the unlit side was recorded. When the mouse crossed over into the unlit compartment, the guillotine door closed and a mild foot shock at 0.5 mV was delivered. The mouse then remained in the dark chamber for ten seconds, a post conditioned stimulus period. Following this, the unlit compartment was illuminated with the house light and the guillotine door was opened. This ensured animals were conditioned to the light/dark cues and not a particular side of the apparatus. Mice were trained to remain in the lit chamber for 120 s. The total number of trials it took to reach criterion was determined for each mouse. On Day 2, mice were placed into the lit compartment and their latency to enter the dark side was recorded.

Experiment 2: The same procedure as stated above was repeated for Experiment 2 except mice were tested on PD72–73, due to equipment availability.

### 2.7. Perfusions

Mice were euthanized on either PD80 (Experiment 1) or PD81 (Experiment 2). First, they were anesthetized with 100 mg/kg sodium pentobarbital *via i.p.* injection and then perfused transcardially with 4% paraformaldehyde in phosphate buffer solution (PBS; 0.287% sodium phosphate monobasic anhydrous, 1.102% sodium phosphate dibasic anhydrous, 0.9% sodium chloride in water). Brains were postfixed overnight and transferred to 30% sucrose in PBS for long-term storage. Brains were sectioned using a cryostat into 40 µm thick coronal sections and stored in cryoprotectant (30% ethylene glycol, 25% glycerin, 45% PBS) at –20 °C.

#### 2.7.1. BrdU-diaminobenzidine (DAB; Experiment 1 and 2)

To detect levels of survival in BrdU-positive (newly divided) cells in the dentate gyrus BrdU-DAB staining was performed. For each mouse, a one-in-six series of sections (*i.e.*, series of sections throughout the rostro-caudal extent of the brain with 240 µm increments separating each section) was stained. Free-floating sections were washed in tris-buffer solution (TBS; 1.3% Trizma hydrochloride, 0.19% Trizma base, 0.9% sodium chloride) and then treated with 0.6% hydrogen peroxide in TBS. To denature DNA, sections were treated with 50% deionized formamide, 10% 20 x saline-sodium citrate (SSC) buffer, 2N hydrochloric acid, and 0.1 M boric acid. Sections were then treated with 0.1% Triton-X and 3% goat serum in TBS (TBS-X plus) and incubated in primary antibody against BrdU made in rat (OBT0030; Accurate, Westbury, NY, USA) at a dilution of 1:200 in TBS-X plus for 72 h at 4 °C. Sections were then washed in TBS, blocked with TBS-X plus for 30 min, and incubated in secondary antibody against rat made in goat (BA-9400; Vector Laboratories, Burlingame CA, USA) at a dilution of 1:250 in TBS-X plus for 100 min at room temperature. Sections were treated using the Vectastain Elite ABC Kit (PK-6200; Vector, Burlingame, CA, USA) and stained using a DAB kit (D4418-505ET; Sigma, St. Louis, MO, USA).

#### 2.7.2. Doublecortin (DCX)-DAB (Experiment 2)

To estimate total numbers of immature neurons in the granule cell layer of the dentate gyrus at the time when the mice were euthanized, DCX-DAB staining was conducted. The immunohistochemistry proceeded similar to BrdU-DAB except omitting the DNA denaturing steps. The primary antibody was goat anti-DCX (1:1000; sc-8066; Santa Cruz Biotechnology, Santa Cruz, CA); secondary

antibody was biotinylated donkey anti-goat (1:200; sc-2042; Santa Cruz Biotechnology, Santa Cruz, CA).

### 2.8. Image analysis

Following Clark and colleagues [12], the entire bilateral granule cell layer, represented in the one-in-six series, was photographed by systematically advancing the field of view of the Zeiss brightfield light microscope and taking multiple photographs, via AxioCam interfaced to computer, under 10 $\times$  (total 100 $\times$ ) magnification. A large depth of field was used so that all particles within the section were visible in each photograph. For BrdU-DAB, these photographs were then analyzed using ImageJ software (NIH, Bethesda, MD, USA) to generate estimates of total number of BrdU-labeled cells per cubic micrometer dentate gyrus sampled and area of the granule cell layer within the sections. Specifically, in each image, the granule cell layer was traced and BrdU-positive nuclei were counted within the traced region automatically by setting a fixed threshold to remove background, and adjusted for double counting at the plane of the section. A subset of animals were also hand counted, and the hand counts regressed against the automated values to establish accurate counts of total numbers of labeled neurons. The R<sup>2</sup> value for these relationships were greater than 90%. The equation was generated from individual sections from multiple individuals representing the full range in cell counts from among the smallest number to the largest number. Hence the equation describes the relationship for all groups and all individuals. Volume of the granule cell layer was estimated by tracing the 2-dimensional area of the granule cell layer in the bilateral 1-in-6 series, and multiplying by the distance between sections (240  $\mu$ m). For DCX-DAB, the protocol mirrored that of BrdU-DAB image analysis except that DCX-positive cells were hand-counted.


### 2.9. Statistical analysis

Data were analyzed using SPSS (version 21) statistical software. In all analyses,  $p < 0.05$  was considered statistically significant. Given that whole litters were injected with either saline or ethanol, data were first analyzed with litter included as a random effect. Litter was not found to be significant on any measure, therefore we removed it for all reported statistics. Thus, the following variables were analyzed using a repeated measure analysis with Day as the within measure and Treatment (alcohol vs. saline) and Intervention (runner vs. sedentary) as the between factors: body weights (g), latency to fall from the Rotarod (s). The following variables were analyzed using a two-way analysis of variance (ANOVA), with Treatment (alcohol vs. saline) and Intervention (runner vs. sedentary) as the two factors: brain weights (g), total number of BrdU+ and DCX+ cells, average distance run on wheels, number of trials to criterion, latency to cross in Passive avoidance (s). Post-hoc analyzes (LSD) were used identify significant pair-wise differences between means ( $p < 0.05$ ).

## 3. Results

### 3.1. Body weights

All mice continued to gain weight throughout treatments (Table 1). Data from the neonatal period (PD5, 7, 9) versus adolescence and adulthood (PD35, 44, 80, 81) were analyzed separately. In Experiment 1, despite litters being randomly assigned to a Treatment, there was a significant main effect of Treatment ( $F_{1,69} = 55.8$ ,  $p < 0.001$ ), wherein alcohol-exposed mice were heavier than saline-exposed mice prior to neonatal treatment. This main effect of Treatment was still evident on PD35 ( $F_{1,34} = 4.2$ ,  $p < 0.05$ ). However, by PD44, alcohol and saline mice had similar body weights and



**Fig. 2.** Wheel Running Levels. Average daily running distance (km) of alcohol-exposed and saline-exposed mice for Experiment 1 and 2. In Experiment 2, saline-exposed mice ran significantly more than alcohol-exposed mice. All values represent mean  $\pm$  SEM. \* $p < 0.05$ .

remained similar at PD80. These data suggest that the single alcohol exposure on PD7 had no long-term impact on body weights.

In Experiment 2, neonatal body weights were collected on PD5, PD7 and PD9. Overall, there was a main effect of Day ( $F_{1,59} = 416.0$ ,  $p < 0.001$ ), a main effect of Treatment ( $F_{1,59} = 11.2$ ,  $p < 0.001$ ) and a Day  $\times$  Treatment interaction ( $F_{1,59} = 22.0$ ,  $p < 0.001$ ). No differences in body weights were evident at the onset of the experiment. However, by PD7, alcohol-exposed mice weighed significantly less than saline-exposed mice ( $p < 0.001$ ). This was still evident on PD9 ( $p < 0.001$ ). A main effect of Treatment remained throughout adolescence (PD35:  $F_{1,33} = 17.4$ ,  $p < 0.001$ ; PD44:  $F_{1,33} = 10.7$ ) and adulthood ( $p < 0.01$ ; PD81:  $F_{1,33} = 11.8$ ,  $p < 0.001$ ), wherein alcohol-exposed mice weighed less than saline-exposed mice. These results suggest that PD5, 7 and 9 produced a long-lasting suppression of growth that persisted to adulthood.

### 3.2. Wheel running

The effects of neonatal ethanol exposure on wheel running behavior depended on the experiment, as indicated by a significant Treatment by Experiment interaction ( $F_{1,29} = 6.6$ ,  $p < 0.01$ ; Fig. 2). In Experiment 1, 5 g/kg ethanol administered on PD7 had no significant influence on wheel running relative to saline controls ( $F_{1,13} = 2.0$ ,  $p = 0.183$ ), whereas in Experiment 2, 5 g/kg administered on PD5, 7 and 9 significantly decreased running ( $F_{1,16} = 5.4$ ,  $p < 0.05$ ). It is likely that the different alcohol administrations produce different FASD profiles. Whereas the single binge produces no significant effect, the repeated-binge produces a long-lasting suppression of motivation for physical activity, as suggested by the decreased running levels.

### 3.3. Blood ethanol concentrations

A dose of 5 g/kg of ethanol, administered by two injections two hours apart, produced significantly high and long-lasting blood ethanol concentration levels that ranged from 300 to 500 mg/dl at time points 3 h, 5 h, and 9 h (Fig. 3). Results of the ANOVA indicated a significant effect of time point. All pairwise comparisons were different ( $p < 0.05$ ) except 9 h vs 30 min, and 5 h vs 3 h. These results are consistent with those of Ikonomidou et al. [35].

### 3.4. Brain weights

In Experiment 2, following perfusion, the brains of mice were weighed in order to determine whether alcohol exposure had long-term effects on brain weight. A t-test revealed

**Table 1**


Effect of postnatal alcohol exposure on body weight (g).

		PD 5	PD 7	PD 9	PD 35	PD 44	PD 80/81
Experiment 1	Saline	N/A	3.02 ± 0.08	N/A	17.37 ± 0.44	18.41 ± 0.37	22.48 ± 0.44
	Alcohol	N/A	3.79 ± 0.07**	N/A	18.42 ± 0.26*	19.06 ± 0.29	23.31 ± 0.42
Experiment 2	Saline	3.02 ± 0.06	4.20 ± 0.10	5.34 ± 0.15	18.20 ± 0.42	18.69 ± 0.49	23.58 ± 0.41
	Alcohol	3.07 ± 0.10	3.58 ± 0.11**	4.55 ± 0.14**	16.20 ± 0.27**	16.77 ± 0.36*	21.80 ± 0.33**

PD, postnatal day. The weights are reported as group means ± SEM.

\* p &lt; 0.05.


\*\* p &lt; 0.001.



**Fig. 3.** Blood Ethanol Concentrations. On PD 7, mice were administered 5 g/kg ethanol (split into two doses, two hours apart). Data points refer to the time after the first of the two injections. The blood ethanol curve illustrates that this dose is sufficient to produce blood ethanol concentrations that exceed the toxic threshold limit (200 mg/dl) for over eight hours.

a main effect of Treatment ( $t_{33}=9.1$ ,  $p < 0.001$ ). The brains of alcohol-exposed mice weighed significantly less than those of saline-exposed mice (Fig. 4A). This decrease was also evident in the calculated brain/body weights ratios for both Treatment groups (Saline:  $0.0188 \text{ g} \pm 0.0004 \text{ g}$ , Alcohol:  $0.0176 \text{ g} \pm 0.0003 \text{ g}$ ). Further, when PD81 body mass was included as a covariate in an ANCOVA, body mass was significantly correlated with brain weight ( $F_{1,31} = 30.6$ ,  $p < 0.001$ ) and the effect of Treatment was also significant ( $F_{1,31} = 51.8$ ,  $p < 0.001$ ). These results suggest that body mass was correlated with brain weight within each group but that matched for body size, alcohol-exposed mice had reduced brain weight as compared to saline-exposed mice (Fig. 4B).

#### Experiment 2: PD 5, 7, 9 Exposure



**Fig. 4.** PD 5, 7 and 9 alcohol exposure produced long-term effects on brain weight as adults. A. The brains of alcohol-exposed mice were significantly smaller than those of saline-exposed mice. B. When matched for body size, alcohol-exposed mice had reduced brain weight as compared to saline-exposed mice. All values represent mean ± SEM. \*p < 0.05.

#### 3.5. BrdU+ cell counts

Exercise increased total number of BrdU+ cells (Fig. 5A, B). ANOVAs revealed a main effect of Intervention in Experiments 1 ( $F_{1,30} = 18.1$ ,  $p < 0.001$ ) and 2 ( $F_{1,30} = 18.1$ ,  $p < 0.001$ ) (Fig. 5D, E). Posthoc comparisons indicated the effect occurred in both alcohol ( $p < 0.01$ ) and saline ( $p < 0.01$ ) mice. No differences in the number of surviving BrdU+ cells between alcohol-exposed and saline-exposed mice was evident in either the running or sedentary mice.


#### Doublecortin (DCX)+ cell counts

Experiment 2: Exercise increased the number of immature neurons (DCX+ cells) in the dentate gyrus granule cell layer (Fig. 5C), while the PD5, 7 and 9 alcohol exposure had no impact. This was indicated by a main effect of Intervention ( $F_{1,20} = 6.4$ ,  $p < 0.05$ ), but no significant effect of Treatment or Treatment x Intervention interaction (Fig. 5F, G).

#### 3.6. Rotarod

Experiment 1: A repeated-measures ANOVA revealed a main effect of Day ( $F_{1,32} = 30.8$ ,  $p < 0.01$ ), signifying improved performance in all mice as the days progressed (Fig. 6A). There was a trend for a main effect of Intervention ( $p = 0.059$ ) and Treatment ( $p = 0.087$ ), but no significant Interaction ( $F < 1$ ).

Experiment 2: A repeated-measures ANOVA revealed, a main effect of Day ( $F_{1,32} = 15.3$ ,  $p < 0.001$ ) along with a Day x Treatment interaction ( $F_{1,32} = 4.3$ ,  $p < 0.05$ ), suggesting saline mice showed greater improvement in performance across days compared to their alcohol counterparts (Fig. 6B). In addition, both a main effect of Treatment ( $F_{1,32} = 6.4$ ,  $p < 0.05$ ), and a main effect of Intervention ( $F_{1,32} = 4.2$ ,  $p < 0.05$ ) were observed, indicating that overall, alcohol



**Fig. 5.** Running increases levels of adult neurogenesis. Total number of bromodeoxyuridine (BrdU) positive cells is used as an estimate of hippocampal neurogenesis. Running significantly increased the number of BrdU+ cells in alcohol-exposed and saline-exposed mice in both Experiment 1 (A) and Experiment 2 (B) as well as the number of DCX+ cells in Experiment 2 (C). Representative image of a BrdU/DAB stained coronal section from a saline-sedentary (D) and saline-runner (E). Representative image of a DCX/DAB stained coronal section of a saline-sedentary (F) and saline-runner (G). Note: differences between experiments in the number of BrdU+ cells are expected due to variation in the BrdU immunohistochemical signal to noise between batches. All values represent mean ± SEM. # $p < 0.001$ , \* $p < 0.05$ .

impaired performance on the Rotarod, while runners exhibited an enhanced performance.


LSD post hoc tests indicate alcohol-sedentary mice displayed worse performance than both saline-sedentary ( $p < 0.05$ ) and saline-runners ( $p < 0.001$ ) but not alcohol-runners. Moreover, alcohol-runners displayed worse performance than saline-runners ( $p < 0.01$ ) but not saline-sedentary mice. These data suggest that a PD5, 7 and 9 alcohol exposure impairs performance on the Rotarod in adulthood, and that running can recover some of the deficit.

### 3.7. Passive avoidance

Experiment 1: On Day 1, no significant differences between groups were observed (Fig. 7A). On Day 2, a main effect of Intervention ( $F_{1,29} = 5.0$ ,  $p < 0.05$ ) was evident (Fig. 7B), wherein all runners remembered the Passive avoidance task better as they had a greater latency to enter the dark chamber than did sedentary mice. How-

ever, neither a Treatment by Intervention nor a main effect of Treatment was found. Together, these data indicate that a single PD7 binge exposure to alcohol does not impact performance on the Passive avoidance task in adulthood, while running enhances performance regardless of neonatal treatment.

Experiment 2: On Day 1, a main effect of Treatment ( $F_{1,33} = 90.0$ ,  $p < 0.01$ ), Intervention ( $F_{1,32} = 67.0$ ,  $p < 0.001$ ) and a Treatment by Intervention interaction ( $F_{1,31} = 60.4$ ,  $p < 0.01$ ) were observed (Fig. 7C). Alcohol-sedentary mice showed a profound deficit in task acquisition, requiring a significantly higher number of trials to avoid entering the dark chamber compared to alcohol-runner ( $p < 0.01$ ), saline-runner ( $p < 0.01$ ) and saline-sedentary mice ( $p < 0.05$ ). On Day 2, a main effect of Intervention ( $F_{1,31} = 5.4$ ,  $p < 0.05$ ) was found wherein runners remembered the Passive avoidance task better as they had a greater latency to enter the dark chamber than did sedentary mice (Fig. 7D). Neither a Treatment x Intervention nor a main effect of Treatment was evident.



**Fig. 6.** Impact of neonatal alcohol exposure on Rotarod performance is dependent on pattern of administration. A. PD 7 alcohol exposure fails to impact Rotarod performance in adulthood on both Days of Rotarod testing. Running wheel access did not improve Rotarod performance in either saline or alcohol-exposed mice. All groups displayed enhanced performance on Day 2 as compared to Day 1. B. PD 5, 7 and 9 alcohol exposure leads to long-term deficits on Rotarod performance, as alcohol-sedentary mice failed to improve across days. Access to a running wheel enhanced Rotarod performance. All values represent mean  $\pm$  SEM. \* $p$  < 0.05.

Overall, these data indicate that PD5, 7, and 9 alcohol exposure is sufficient to impact acquisition but not retention of the Passive avoidance task in adulthood, while running enhances retention of the task regardless of neonatal treatment.

#### 4. Discussion

These results demonstrate that a third-trimester equivalent exposure to either a single or multiple binges of alcohol has no long-lasting influence on the number of surviving BrdU+ cells in the C57BL/6J mouse model (Fig. 5). This is important given that previous literature was mixed on the issue. It is possible that the effect of neonatal alcohol on the number of surviving BrdU+ cells is strain dependent. Further, a single binge was not sufficient, while multiple binges were sufficient, to cause observable behavioral deficits in Passive avoidance and Rotarod performance (Figs. 6, 7). These data support the use of a three-binge model in C57BL/6J mice for inducing reliable behavioral deficits on these tasks. Finally, the data support the idea that exercise can ameliorate alcohol-induced cognitive deficits, and that this occurs in concert with increased adult hippocampal neurogenesis. However, because neonatal alcohol exposure did not impair adult hippocampal neurogenesis, these results suggest that effects of exercise provide global compensation rather than repair specific damages induced from neonatal alcohol exposure.




Results of our study suggest that effects of neonatal alcohol exposure on the number of surviving BrdU+ cells and immature neurons (DCX+ cells) may be much smaller and dependent on methodological differences than originally hypothesized. The

vast majority of surviving cells in the hippocampal dentate gyrus are neurons [11,30,32,65,66], while less than 10% are glial cells [13,16,25]. Research suggests that these percentages are not influenced by developmental alcohol exposure [9,30]. Therefore, we conclude that neonatal alcohol exposure (both single and multiple binges) did not impact levels of adult hippocampal neurogenesis (Fig. 5). These data are consistent with a growing number of studies in rodents [21,30,42,55,71]. However, other studies implicate neonatal alcohol exposure as the cause of reduced cell survival [24,26,27,34,42]. We now interpret these differences as likely related to strain, age or method of measuring neurogenesis. No effect was observed in C57BL/6J mice [71], while CD1 mice exhibited a deficit [34]. Prenatal ethanol significantly decreased levels of adult hippocampal neurogenesis in adult but not adolescent rats [22]. In studies from the same lab, multiple injections [30] versus a single injection of BrdU [24,26,27] produce differing results. Together, these data suggest that the impact of neonatal alcohol exposure on levels of adult hippocampal neurogenesis may not be as robust as originally hypothesized and the differences in the literature could simply be due to idiosyncratic methodological differences.

In rodent models of FASD, different binge schedules influence behavioral outcome and we wanted to determine the appropriate binge schedule for inducing reliable behavioral deficits on the Passive avoidance and Rotarod tasks, because each of these tasks involves a different subset of vulnerable neuronal substrates. What we discovered was that a multiple-binge produced behavioral deficits in both tasks whereas the single binge had no effect (Fig. 6B; 7A, C). Other multiple-binge paradigms have been shown to impair performance on Trace conditioning [52], Morris water maze [23] and Rotarod [40] in rodents. Wagner et al. [68] demonstrated a single PD7 binge alcohol exposure produced learning deficits in PD70-79 using the Morris water maze task in C57BL/6J mice. However, a previous study found this only to be true in adolescents not adults [71]. It is possible that the Morris water maze involves a different subset of neuronal substrates than either the Passive avoidance or Rotarod tasks do individually. In any case, results from our study suggest that the three-binge model is better suited for detecting behavioral deficits on the Passive avoidance and Rotarod tasks.

The Passive avoidance task is useful in its design: to successfully perform the task, the animal must learn to not move. It does not require the animal to be able to swim successfully or run smoothly, but rather the animal just needs to stay in the same place. Therefore, it could be hypothesized that animals with impaired motor coordination would be, by default, good at this task. Yet, our results show that despite an impaired performance on the Rotarod task, PD 5, 7 and 9 alcohol-exposed mice require a higher number of trials to reach criterion on the Passive avoidance task (*i.e.* exhibit increased motor activity). This suggests, that the behavioral deficits exhibited on the Passive avoidance task are not the result of any motor impairment but rather are consistent with the hallmark FASD features of increased hyperactivity and attention deficits.

Voluntary wheel running was an effective intervention. Not only did it increase adult neurogenesis (Fig. 5) and improve motor coordination on the Rotarod (Fig. 6B), but it also enhanced performance on the Passive avoidance task (Fig. 7). The beneficial influence of voluntary exercise has been well established in the healthy brain. Motor coordination and learning on the Rotarod are likely related to exercise-induced changes in the cerebellum [15,36]. Passive avoidance requires an animal to learn and remember an association between a behavioral action, moving to the dark side of the chamber, with an aversive stimulus, the electric shock. This type of spatial and associative learning and memory involves the hippocampus, and potentially hippocampal neurogenesis [65]. However, if increased neurogenesis was involved, our data suggest

**Experiment 1: PD 7 Exposure****Experiment 2: PD 5, 7, 9 Exposure**

**Fig. 7.** PD 5, 7 and 9 administration but not PD 7 alone impairs Passive avoidance performance. A. Number of trials to reach criterion on Day 1 for Experiment 1. There were no differences between alcohol-exposed and saline-exposed mice as well as no differences between running and sedentary mice. B. Latency to cross over into the dark chamber on Day 2 for Experiment 1. Exercise significantly increased the latency to cross over for both the alcohol-exposed and the saline exposed-mice. C. Number of trials to learn the Passive avoidance task on Day 1 for Experiment 2. Alcohol exposure significantly increased the number of trials it took to reach criterion for sedentary mice. Running significantly decreased the trials to criterion for alcohol-exposed mice. D. Latency to cross over into the dark chamber on Day 2 for Experiment 2. Exercise significantly increased the latency to cross over in both the alcohol-exposed and the saline-exposed mice. All values represent mean  $\pm$  SEM. \* $p < 0.05$ , # $p < 0.001$ .

the effect was through compensation rather than repair because the alcohol-exposed mice did not show impaired neurogenesis.

Alternatively, behavioral improvements from exercise in our study may have nothing to do with neurogenesis. Exercise upregulates a number of trophic factors throughout the brain [15,59,67]. These trophic factors promote the differentiation, neurite extension and survival of a variety of neuronal populations in many different brain regions, not only the hippocampus [13,15,59]. Exercise promotes cerebellar plasticity. In mice raised in an environment with continual access to free running the dendritic trees of cerebellar Purkinje cells are significantly more complex and contain a greater number of spines when compared to those of their control littermates [57]. Further, complex motor skill learning has been associated with increased numbers of parallel fiber to Purkinje cell synapses [39] as well as increased BDNF expression in the cerebellum [41]. Therefore, it is likely that exercise ameliorated deficits in the Rotarod task by affecting plasticity in the cerebellum. Similarly, Passive avoidance performance could have been improved *via* changes in the prefrontal cortex [38,69]. In addition to learning and remembering the association, Passive avoidance requires the mouse to suppress the innate desire to cross over to the dark side, and a prominent brain region involved in behavioral inhibition is the prefrontal cortex. Third-trimester equivalent alcohol exposure is known to impact prefrontal anatomy [25,28,45]. Prenatal alcohol exposure induced both significant cell loss in the prefrontal cortex and impaired reversal learning on a go/no-go procedure in rats [50]. Additionally, heavy prenatal alcohol expo-

sure impaired performance on response-inhibition tasks in humans [19,70]. More recently, exercise has been shown to partially reverse neonatal alcohol-induced deficits in the morphology of prefrontal Layer II/III pyramidal neurons [25]. Therefore, it is possible that exercise rescued Passive avoidance deficits in our study by repairing morphological deficits in multiple brain regions.

It is important to acknowledge certain limitations within this study. First, the model itself brings with it certain caveats. Undoubtedly, there is a mild form of neonatal stress through the alcohol administration. However, to our knowledge, this is true of all the models. In the commonly used pair-feeding method, alcohol-treated dams eat *ad libitum*, while their pair-fed counterparts are calorie restricted, thus introducing a stress component [31]. Further gastric intubation of pups is inherently stressful, given the inclusion of the sham-intubated group, which serves as a stress control. Moreover, the gastric intubation is typically done in rats, it is exceedingly challenging in mice given the small size of the pups. Thus, there does not appear to be one model that is one hundred percent stress free. Second, given the need to record individual running levels, mice were singly housed. This brings with it certain caveats as studies have shown that single housing can induce stress responses in both rats [8] and mice [5]. For example, single housing during adolescence not only reduced the explorative behavior of CD-1 female mice but also their locomotor activity was increased [29]. Further, single housing in C57BL/6J adult male mice increased anxiety and depressive-like behaviors and also reduced BDNF levels, notably in the frontal cortex [5]. However, a growing literature

suggests that single housing may not be stressful for mice. Studies demonstrate that single housing does not increase stress markers in male mice compared with group housing [2,33]. In fact, one study found that group-housed male mice actually exhibited significantly higher corticosterone levels than did their single-housed counterparts [37]. Overall, the literature suggests that the impact of single housing is not ideal, but important nonetheless in this animal model, given the need to record individual running levels. Third, clearly the PD 5, 7 and 9 alcohol-exposed pups suffered some undernutrition during the alcohol exposure procedure given the short-term reduction in body weight, which in this case extended all the way into adulthood. It is possible that the short-term under nourishment contributed to the long lasting effects observed for body mass, brain mass and behavior. Under nourishment would have occurred for both temporary loss of macro and micronutrients during a period of massive growth. One well-studied essential nutrient during development is choline. Choline-deficient pups exhibit impaired learning and memory when tested later in life [18,51,56]. Interestingly, choline supplementation consistently has been shown to attenuate behavioral deficits in rat pups exposed to fetal alcohol [62–64,72]. Therefore, some of the deficits we observed from our neonatal alcohol treatment could have exerted their influence through under nourishment rather than pharmacological effects directly related to ethanol. This is not necessarily a limitation of the model but an important consideration since many fetal alcohol babies are born underweight [44,48,49], and the low birth weights may be attributed in part to the moms getting most of their calories from alcohol [49].

## 5. Conclusion

Together, these results illustrate that PD 5, 7 and 9 alcohol exposure causes behavioral deficits on both the Passive avoidance and the Rotarod task while having no impact on adult hippocampal neurogenesis levels in the C57BL/6J mouse model. Passive avoidance is a unique behavioral task that can assess multiple neural substrates, while requiring minimum strenuous activity from the animal to ensure the results do not simply reflect general motor impairments. In the current study, despite performing poorly on the Rotarod task, the alcohol-exposed animals required significantly more trials to learn the Passive Avoidance task, suggesting this was a learning not a motor deficit. In addition, these data illustrate the potential utility of exercise as a therapeutic intervention for both fetal alcohol-induced motor and cognitive deficits. It is possible that the beneficial effects of exercise on behavioral performance could be partially mediated by increased neurogenesis via compensation but not repair because neurogenesis was not impaired from alcohol. However, it is well established that exercise has many effects on the healthy and diseased brain. Hence, future research should explore alternative mechanisms (i.e. potential repair of prefrontal cortex and cerebellar deficits) as mechanisms underlying behavioral recovery from early postnatal exposure to multiple alcohol binges.

## Funding

This research was supported by Justin Rhodes' lab at the University of Illinois at Urbana-Champaign and funding from NIH 1 RO1 MH083807 (JSR) and NIH 1 RO1 DA027487 (JSR). Additionally, support was provided by the Arnold O. and Mabel M. Beckman Foundation through the Beckman Institute Postdoctoral Fellowship Program at the University of Illinois at Urbana-Champaign (GFH).

## Acknowledgements

This research was supported by Justin Rhodes' lab at the University of Illinois at Urbana-Champaign and funding from NIH 1 RO1 MH083807 (JSR) and NIH 1 RO1 DA027487 (JSR). Additionally, support was provided by the Arnold O. and Mabel M. Beckman Foundation through the Beckman Institute Postdoctoral Fellowship Program at the University of Illinois at Urbana-Champaign (GFH). The authors would like to extend their sincere gratitude to the Beckman Institute Animal Facility staff for excellent animal care.

## References

- [1] E.L. Abel, In utero alcohol exposure and developmental delay of response inhibition, *Alcohol Clin. Exp. Res.* 6 (3) (1982) 369–376.
- [2] S.S. Arndt, M.C. Laarakker, H.A. van Lith, F.J. van der Staay, E. Gieling, A.R. Salomons, J. van't Klooster, F. Ohl, Individual housing of mice—impact on behaviour and stress responses, *Physiol. Behav.* 97 (3–4) (2009) 385–393.
- [3] S. Barron, E.P. Riley, Passive avoidance performance following neonatal alcohol exposure, *Neurotoxicol. Teratol.* 12 (2) (1990) 135–138.
- [4] H.C. Becker, C.L. Randall, Effects of prenatal ethanol exposure in C57BL mice on locomotor activity and passive avoidance behavior, *Psychopharmacology (Berl.)* 97 (1) (1989) 40–44.
- [5] A. Berry, V. Bellisario, S. Capoccia, P. Tirassa, A. Calza, E. Alleva, F. Cirulli, Social deprivation stress is a triggering factor for the emergence of anxiety- and depression-like behaviours and leads to reduced brain BDNF levels in C57BL/6J mice, *Psychoneuroendocrinology* 37 (6) (2012) 762–772.
- [6] J. Brown, C.M. Cooper-Kuhn, G. Kempermann, H. Van Praag, J. Winkler, F.H. Gage, H.G. Kuhn, Enriched environment and physical activity stimulate hippocampal but not olfactory bulb neurogenesis, *Eur. J. Neurosci.* 17 (10) (2003) 2042–2046.
- [7] F. Calhoun, M.L. Attilia, P.A. Spagnolo, C. Rotondo, R. Mancinelli, M. Ceccanti, National Institute on Alcohol Abuse and Alcoholism and the study of fetal alcohol spectrum disorders. The International Consortium, *Ann. Ist Super Sanita* 42 (1) (2006) 4–7.
- [8] A.M. Chappell, E. Carter, B.A. McCool, J.L. Weiner, Adolescent rearing conditions influence the relationship between initial anxiety-like behavior and ethanol drinking in male Long Evans rats, *Alcohol Clin. Exp. Res.* 37 (Suppl. (1)) (2013) E394–E403.
- [9] I.Y. Choi, A.M. Allan, L.A. Cunningham, Moderate fetal alcohol exposure impairs the neurogenic response to an enriched environment in adult mice, *Alcohol Clin. Exp. Res.* 29 (11) (2005) 2053–2062.
- [10] B.R. Christie, S.E. Swann, C.J. Fox, D. Froc, S.E. Lieblich, V. Redila, A. Webber, Voluntary exercise rescues deficits in spatial memory and long-term potentiation in prenatal ethanol-exposed male rats, *Eur. J. Neurosci.* 21 (6) (2005) 1719–1726.
- [11] P.J. Clark, W.J. Brzezinska, E.K. Puchalski, D.A. Krone, J.S. Rhodes, Functional analysis of neurovascular adaptations to exercise in the dentate gyrus of young adult mice associated with cognitive gain, *Hippocampus* 19 (10) (2009) 937–950.
- [12] P.J. Clark, W.J. Brzezinska, M.W. Thomas, N.A. Ryzhenko, S.A. Toshkov, J.S. Rhodes, Intact neurogenesis is required for benefits of exercise on spatial memory but not motor performance or contextual fear conditioning in C57BL/6 mice, *Neuroscience* 155 (4) (2008) 1048–1058.
- [13] S.J. Colcombe, A.F. Kramer, K.I. Erickson, P. Scalf, E. McAuley, N.J. Cohen, A. Webb, G.J. Jerome, D.X. Marquez, S. Elavsky, Cardiovascular fitness, cortical plasticity, and aging, *Proc. Natl. Acad. Sci. U. S. A.* 101 (9) (2004) 3316–3321.
- [14] P.D. Connor, P.D. Sampson, F.L. Bookstein, H.M. Barr, A.P. Streissguth, Direct and indirect effects of prenatal alcohol damage on executive function, *Dev. Neuropsychol.* 18 (3) (2000) 331–354.
- [15] C.W. Cotman, N.C. Berchtold, Exercise: a behavioral intervention to enhance brain health and plasticity, *Trends Neurosci.* 25 (6) (2002) 295–301.
- [16] F.T. Crews, A. Mdzinarishvili, D. Kim, J. He, K. Nixon, Neurogenesis in adolescent brain is potently inhibited by ethanol, *Neuroscience* 137 (2) (2006) 437–445.
- [17] J. Dobbing, J. Sands, Comparative aspects of the brain growth spurt, *Early Hum. Dev.* 3 (1) (1979) 79–83.
- [18] D.A. Fernandes de Abreu, E. Nivet, N. Baril, M. Khrestchatsky, F. Roman, F. Feron, Developmental vitamin D deficiency alters learning in C57Bl/6J mice, *Behav. Brain Res.* 208 (2) (2010) 603–608.
- [19] S.L. Fryer, S.F. Tapert, S.N. Mattson, M.P. Paulus, A.D. Spadoni, E.P. Riley, Prenatal alcohol exposure affects frontal-striatal BOLD response during inhibitory control, *Alcohol Clin. Exp. Res.* 31 (8) (2007) 1415–1424.
- [20] J. Gil-Mohapel, F. Boehme, L. Kainer, B.R. Christie, Hippocampal cell loss and neurogenesis after fetal alcohol exposure: insights from different rodent models, *Brain Res. Rev.* 64 (2) (2010) 283–303.
- [21] J. Gil-Mohapel, F. Boehme, A. Patten, A. Cox, L. Kainer, E. Giles, P.S. Brocardo, B.R. Christie, Altered adult hippocampal neuronal maturation in a rat model of fetal alcohol syndrome, *Brain Res.* 1384 (2011) 29–41.
- [22] J. Gil-Mohapel, A.K. Titterness, A.R. Patten, S. Taylor, A. Ratzlaff, T. Ratzlaff, J. Helfer, B.R. Christie, Prenatal ethanol exposure differentially affects

- hippocampal neurogenesis in the adolescent and aged brain, *Neuroscience* 273 (2014) 174–188.
- [23] C.R. Goodlett, T.B. Johnson, Neonatal binge ethanol exposure using intubation: timing and dose effects on place learning, *Neurotoxicol. Teratol.* 19 (6) (1997) 435–446.
- [24] G.F. Hamilton, K.E. Boschen, C.R. Goodlett, W.T. Greenough, A.Y. Klintsova, Housing in environmental complexity following wheel running augments survival of newly generated hippocampal neurons in a rat model of binge alcohol exposure during the third trimester equivalent, *Alcohol Clin. Exp. Res.* 36 (7) (2012) 1196–1204.
- [25] G.F. Hamilton, K.J. Criss, A.Y. Klintsova, Voluntary exercise partially reverses neonatal alcohol-induced deficits in mPFC layer II/III dendritic morphology of male adolescent rats, *Synapse* 69 (8) (2015) 405–415.
- [26] G.F. Hamilton, S.A. Jablonski, F.L. Schiffino, S.A. St Cyr, M.E. Stanton, A.Y. Klintsova, Exercise and environment as an intervention for neonatal alcohol effects on hippocampal adult neurogenesis and learning, *Neuroscience* 265 (2014) 274–290.
- [27] G.F. Hamilton, N.J. Murawski, S.A. St Cyr, S.A. Jablonski, F.L. Schiffino, M.E. Stanton, A.Y. Klintsova, Neonatal alcohol exposure disrupts hippocampal neurogenesis and contextual fear conditioning in adult rats, *Brain Res.* 1412 (2011) 88–101.
- [28] G.F. Hamilton, L.T. Whitcher, A.Y. Klintsova, Postnatal binge-like alcohol exposure decreases dendritic complexity while increasing the density of mature spines in mPFC Layer II/III pyramidal neurons, *Synapse* 64 (2) (2010) 127–135.
- [29] M. Haupt, A.T. Schaefers, Effects of postweaning social and physical deprivation on locomotor activity patterns and explorative behavior in female CD-1 mice, *Dev. Psychobiol.* 52 (4) (2010) 383–393.
- [30] J.L. Helfer, C.R. Goodlett, W.T. Greenough, A.Y. Klintsova, The effects of exercise on adolescent hippocampal neurogenesis in a rat model of binge alcohol exposure during the brain growth spurt, *Brain Res.* 1294 (2009) 1–11.
- [31] K.G. Hellemans, J.H. Sliwowska, P. Verma, J. Weinberg, Prenatal alcohol exposure: fetal programming and later life vulnerability to stress, depression and anxiety disorders, *Neurosci. Biobehav. Rev.* 34 (6) (2010) 791–807.
- [32] M.M. Holmes, L.A. Galea, R.E. Mistlberger, G. Kempermann, Adult hippocampal neurogenesis and voluntary running activity: circadian and dose-dependent effects, *J. Neurosci. Res.* 76 (2) (2004) 216–222.
- [33] C. Hunt, C. Hambley, Faecal corticosterone concentrations indicate that separately housed male mice are not more stressed than group housed males, *Physiol. Behav.* 87 (3) (2006) 519–526.
- [34] A. Ieraci, D.G. Herrera, Single alcohol exposure in early life damages hippocampal stem/progenitor cells and reduces adult neurogenesis, *Neurobiol. Dis.* 26 (3) (2007) 597–605.
- [35] C. Ikonomidou, P. Bittigau, M.J. Ishimaru, D.F. Wozniak, C. Koch, K. Genz, M.T. Price, V. Stefovská, F. Horster, T. Tenkova, et al., Ethanol-induced apoptotic neurodegeneration and fetal alcohol syndrome, *Science* 287 (5455) (2000) 1056–1060.
- [36] K.R. Isaacs, B.J. Anderson, A.A. Alcantara, J.E. Black, W.T. Greenough, Exercise and the brain: angiogenesis in the adult rat cerebellum after vigorous physical activity and motor skill learning, *J. Cereb. Blood Flow Metab.* 12 (1) (1992) 110–119.
- [37] R. Kamakura, M. Kovalainen, J. Leppaluoto, K.H. Herzig, K.A. Makela, The effects of group and single housing and automated animal monitoring on urinary corticosterone levels in male C57BL/6 mice, *Physiol. Rep.* 4 (3) (2016).
- [38] E.D. Kemble, J.T. Tapp, Passive and active avoidance performance following small amygdaloid lesions in rats, *Physiol. Behav.* 3 (5) (1968) 713–.
- [39] J.A. Kleim, R.A. Swain, K.A. Armstrong, R.M. Napper, T.A. Jones, W.T. Greenough, Selective synaptic plasticity within the cerebellar cortex following complex motor skill learning, *Neurobiol. Learn. Mem.* 69 (3) (1998) 274–289.
- [40] A.Y. Klintsova, R.M. Cowell, R.A. Swain, R.M. Napper, C.R. Goodlett, W.T. Greenough, Therapeutic effects of complex motor training on motor performance deficits induced by neonatal binge-like alcohol exposure in rats. I. Behavioral results, *Brain Res.* 800 (1) (1998) 48–61.
- [41] A.Y. Klintsova, E. Dickson, R. Yoshida, W.T. Greenough, Altered expression of BDNF and its high-affinity receptor TrkB in response to complex motor learning and moderate exercise, *Brain Res.* 1028 (1) (2004) 92–104.
- [42] A.Y. Klintsova, J.L. Helfer, L.H. Calizo, W.K. Dong, C.R. Goodlett, W.T. Greenough, Persistent impairment of hippocampal neurogenesis in young adult rats following early postnatal alcohol exposure, *Alcohol Clin. Exp. Res.* 31 (12) (2007) 2073–2082.
- [43] C.L. Lantz, W. Wang, A.E. Medina, Early alcohol exposure disrupts visual cortex plasticity in mice, *Int. J. Dev. Neurosci.* 30 (5) (2012) 351–357.
- [44] B. Larroque, M. Kaminski, N. Lelong, D. Subtil, P. Dehaene, Effects of birth weight of alcohol and caffeine consumption during pregnancy, *Am. J. Epidemiol.* 137 (9) (1993) 941–950.
- [45] R.C. Lawrence, N.K. Otero, S.J. Kelly, Selective effects of perinatal ethanol exposure in medial prefrontal cortex and nucleus accumbens, *Neurotoxicol. Teratol.* 34 (1) (2012) 128–135.
- [46] D.J. Livy, E.K. Miller, S.E. Maier, J.R. West, Fetal alcohol exposure and temporal vulnerability: effects of binge-like alcohol exposure on the developing rat hippocampus, *Neurotoxicol. Teratol.* 25 (4) (2003) 447–458.
- [47] A.M. Magarinos, B.S. McEwen, M. Sabouret, P. Pevet, Rapid and reversible changes in intrahippocampal connectivity during the course of hibernation in European hamsters, *Proc. Natl. Acad. Sci. U. S. A.* 103 (49) (2006) 18775–18780.
- [48] P.A. May, J.P. Gossage, W.O. Kalberg, L.K. Robinson, D. Buckley, M. Manning, H.E. Hoyme, Prevalence and epidemiologic characteristics of FASD from various research methods with an emphasis on recent in-school studies, *Dev. Disabil. Res. Rev.* 15 (3) (2009) 176–192.
- [49] P.A. May, K.J. Hamrick, K.D. Corbin, J.M. Hasken, A.S. Marais, L.E. Brooke, J. Blankenship, H.E. Hoyme, J.P. Gossage, Dietary intake, nutrition, and fetal alcohol spectrum disorders in the Western Cape Province of South Africa, *Reprod. Toxicol.* 46 (2014) 31–39.
- [50] S.M. Mihalick, J.E. Crandall, J.C. Langlois, J.D. Krienke, W.V. Dube, Prenatal ethanol exposure, generalized learning impairment, and medial prefrontal cortical deficits in rats, *Neurotoxicol. Teratol.* 23 (5) (2001) 453–462.
- [51] H.C. Moreno, I. de Brugada, D. Carias, M. Gallo, Long-lasting effects of prenatal dietary choline availability on object recognition memory ability in adult rats, *Nutr. Neurosci.* 16 (6) (2013) 269–274.
- [52] N.J. Murawski, S.A. Jablonski, K.L. Brown, M.E. Stanton, Effects of neonatal alcohol dose and exposure window on long delay and trace eyeblink conditioning in juvenile rats, *Behav. Brain Res.* 236 (1) (2013) 307–318.
- [53] A.K. Olson, B.D. Eadie, C. Ernst, B.R. Christie, Environmental enrichment and voluntary exercise massively increase neurogenesis in the adult hippocampus via dissociable pathways, *Hippocampus* 16 (3) (2006) 250–260.
- [54] A.R. Patten, C.J. Fontaine, B.R. Christie, A comparison of the different animal models of fetal alcohol spectrum disorders and their use in studying complex behaviors, *Front. Pediatr.* 2 (2014) 93.
- [55] A.R. Patten, D.J. Moller, J. Graham, J. Gil-Mohapel, B.R. Christie, Liquid diets reduce cell proliferation but not neurogenesis in the adult rat hippocampus, *Neuroscience* 254 (2013) 173–184.
- [56] J.N. Peak, K.M. Turner, T.H. Burne, The effect of developmental vitamin D deficiency in male and female Sprague-Dawley rats on decision-making using a rodent gambling task, *Physiol. Behav.* 138 (2015) 319–324.
- [57] J.J. Pysh, G.M. Weiss, Exercise during development induces an increase in Purkinje cell dendrite tree size, *Science* 206 (4415) (1979) 230–232.
- [58] C. Rasmussen, Executive functioning and working memory in fetal alcohol spectrum disorder, *Alcohol Clin. Exp. Res.* 29 (8) (2005) 1359–1367.
- [59] P. Rasmussen, P. Brassard, H. Adser, M.V. Pedersen, L. Leick, E. Hart, N.H. Secher, B.K. Pedersen, H. Pilegaard, Evidence for a release of brain-derived neurotrophic factor from the brain during exercise, *Exp. Physiol.* 94 (10) (2009) 1062–1069.
- [60] V.A. Redila, A.K. Olson, S.E. Swann, G. Mohades, A.J. Webber, J. Weinberg, B.R. Christie, Hippocampal cell proliferation is reduced following prenatal ethanol exposure but can be rescued with voluntary exercise, *Hippocampus* 16 (3) (2006) 305–311.
- [61] E.P. Riley, E.A. Lochry, N.R. Shapiro, Lack of response inhibition in rats prenatally exposed to alcohol, *Psychopharmacology (Berl.)* 62 (1) (1979) 47–52.
- [62] S.H. Ryan, J.K. Williams, J.D. Thomas, Choline supplementation attenuates learning deficits associated with neonatal alcohol exposure in the rat: effects of varying the timing of choline administration, *Brain Res.* 1237 (2008) 91–100.
- [63] R.D. Schneider, J.D. Thomas, Adolescent choline supplementation attenuates working memory deficits in rats exposed to alcohol during the third trimester equivalent, *Alcohol Clin. Exp. Res.* 40 (4) (2016) 897–905.
- [64] J.D. Thomas, T.D. Tran, Choline supplementation mitigates trace, but not delay, eyeblink conditioning deficits in rats exposed to alcohol during development, *Hippocampus* 22 (3) (2012) 619–630.
- [65] H. van Praag, B.R. Christie, T.J. Sejnowski, F.H. Gage, Running enhances neurogenesis, learning, and long-term potentiation in mice, *Proc. Natl. Acad. Sci. U. S. A.* 96 (23) (1999) 13427–13431.
- [66] H. van Praag, T. Shubert, C. Zhao, F.H. Gage, Exercise enhances learning and hippocampal neurogenesis in aged mice, *J. Neurosci.* 25 (38) (2005) 8680–8685.
- [67] S. Vaynman, F. Gomez-Pinilla, License to run: exercise impacts functional plasticity in the intact and injured central nervous system by using neurotrophins, *Neurorehabil. Neural. Repair* 19 (4) (2005) 283–295.
- [68] J.L. Wagner, F.C. Zhou, C.R. Goodlett, Effects of one- and three-day binge alcohol exposure in neonatal C57BL/6 mice on spatial learning and memory in adolescence and adulthood, *Alcohol* 48 (2) (2014) 99–111.
- [69] T.J. Walsh, H.A. Tilson, D.L. DeHaven, R.B. Mailman, A. Fisher, I. Hanin, AF64A, a cholinergic neurotoxin, selectively depletes acetylcholine in hippocampus and cortex, and produces long-term passive avoidance and radial-arm maze deficits in the rat, *Brain Res.* 321 (1) (1984) 91–102.
- [70] A.L. Ware, M.A. Infante, J.W. O'Brien, S.F. Tapert, K.L. Jones, E.P. Riley, S.N. Mattson, An fMRI study of behavioral response inhibition in adolescents with and without histories of heavy prenatal alcohol exposure, *Behav. Brain Res.* 278 (2015) 137–146.
- [71] D.F. Wozniak, R.E. Hartman, M.P. Boyle, S.K. Vogt, A.R. Brooks, T. Tenkova, C. Young, J.W. Olney, L.J. Muglia, Apoptotic neurodegeneration induced by ethanol in neonatal mice is associated with profound learning/memory deficits in juveniles followed by progressive functional recovery in adults, *Neurobiol. Dis.* 17 (3) (2004) 403–414.
- [72] C.H. Zhu, T. Wu, Y. Jin, B.X. Huang, R.F. Zhou, Y.Q. Wang, X.L. Luo, H.L. Zhu, Prenatal choline supplementation attenuates spatial learning deficits of offspring rats exposed to low-protein diet during fetal period, *J. Nutr. Biochem.* 32 (2016) 163–170.